

Thermal energy requirement for tobacco caterpillar *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae) development on artificial diet

SUBHASH CHANDER, MAZHAR HUSAIN and GUNDAPPA BARADEVANAL^{1*}

Division of Entomology, ICAR-IARI, New Delhi, India

¹Division of Crop Protection, ICAR-CISH, Lucknow, India

*Corresponding author: Gundappa@icar.gov.in

ABSTRACT

Development and survival of tobacco caterpillar *Spodoptera litura* larvae and pupae were studied at each of seven constant temperatures, 15±1, 18±1, 21±1, 24±1, 27±1, 30±1 and 33±1°C, in incubators. Oviposition, incubation, larval and pupal periods were recorded at each of the temperatures. The survival from 1st instar to adult emergence ranged between a maximum of 60% at 24°C and 30% at each of 18°C and 33°C. Developmental duration of different larval instars and pupae declined with an increase in temperature from 15 to 33°C. The optimum temperature for larval and pupal survival in *S. litura* on artificial diet was observed to be 24°C. On the other hand, optimum temperature range for oviposition was observed to be 27-33°C. The relationship between temperature (T) and development rate (r) was established using linear regression to estimate the thermal constant (K) and development threshold (T₀). Thermal constant for 1st to 6th larval instars were computed as 76.9, 90.9, 125, 83.3, 62.5, 90.9 and 200 degree days (DD), respectively, with corresponding development thresholds as 12.8, 11.7, 9.8, 10.3, 10, 10.9 and 13.2°C. Single developmental threshold for larval stage was computed as 13°C. These thermal constants and development thresholds can be used in developing mechanistic population simulation model, which in turn will facilitate the assessment of climate change impact on the pest.

Key words: Thermal energy, artificial diet, development rate, *Spodoptera litura*, survival

Tobacco caterpillar, *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae) is a perilous insect pest affecting economically important crops such as tobacco (*Nicotiana glauca* L.), castor (*Ricinus communis* L.), cotton (*Gossypium* sp. L.), soybean (*Glycine max* L.) and groundnut (*Arachis hypogea* L.) throughout tropical and temperate Asia, Australasia and the Pacific Islands (EPPO, 2020). Out of 112 globally recorded host plants of *S. litura*, 60 are known only from India (CABI, 2020; Jha *et al.*, 2016). Due to nocturnal habit, high mobility of adult moths and ability to oviposit on a wide range of host plants, *S. litura* has huge potential to invade new areas and to adapt to wide range of ecological situations (Duraimurugan, 2018). In India, *S. litura* is widespread in almost all the states and inflicts significant losses to economically important crops like soybean, cotton and groundnut. *S. litura* is likely to further increase under the imminent climate change, as it has been found to consume 30 per cent more cotton leaves at elevated CO₂ levels than ambient (Kranthi *et al.*, 2009). Outbreaks of *S. litura* were also noticed in major sunflower growing areas of Central and Southern India. During 2005, the outbreak of *S. litura* led to more than 90 per cent defoliation of sunflower (Sujatha and Lakshminarayana, 2007).

Insects being poikilothermic organisms, the developmental rate is highly reliant on external temperature conditions. Hence, temperature is generally considered the single most significant environmental factor influencing behavior, distribution, development, survival and reproduction in insects (Bale *et al.*, 2002). Knowledge on the temperature-dependent population growth potential of insect pests is highly imperative for understanding their population dynamics and implementing agro-eco region specific pest management strategies, especially in the context of predicted global climate warming (Kroschel *et al.*, 2013). Considering this, the temperature rise of 2.7–4.7°C predicted due to potential climate change (IPCC, 2013) may have drastic consequences for future incidence of *S. litura*. The vast majority of studies that infer the effects of temperature on developmental biology of *S. litura* have been undertaken under only one constant temperature in laboratory (Roa *et al.*, 1989; Prasad 2004; Bharathi *et al.*, 2007; Patil *et al.*, 2014). A few studies that addressed the development of *S. litura* at a range of constant temperatures, were concerned with predicting developmental rates and threshold temperatures using linear degree day or heat summation models (Rao *et al.*, 1989). The objective of this study was to develop thermal

constants and developmental thresholds for *S. litura*, which can be used in developing mechanistic population simulation model, which in turn will facilitate the assessment of climate change impact on the pest.

MATERIAL AND METHODS

The study was conducted in the laboratory of the Division of Entomology, ICAR-Indian Agricultural Research Institute (IARI), New Delhi. The laboratory experiments were conducted using Biological Oxygen Demand (BOD) incubators that were set at constant relative humidity ($65 \pm 5\%$) and photoperiod (14:10 L:D) and different temperatures as required for determining thermal constants and development thresholds study of *S. litura*.

Establishment of a stock culture of S. litura

The *S. litura* larvae were collected from IARI fields, sorted out and reared in glass containers on fresh artificial diet that was changed daily to maintain hygienic conditions. The containers' tops were covered with perforated aluminum foil lids. The pupae were kept in separate 20x10 cm rearing jars for adult emergence. On emergence, adults were placed in 20x10 cm rearing jars with their tops covered with muslin cloth and were provided with 15% sugar solution through pieces of cotton in petri dishes. The adult food was daily changed. Fresh tomato branches dipped in 100 ml water containers were placed in adult rearing jars for oviposition. The adult rearing jars were maintained at $25 \pm 1^\circ\text{C}$ and 65% relative humidity. The faecal matter and unused diet was removed from containers as and when required.

Development and survival of S. litura at different constant temperatures

Development and survival of *S. litura* life stages viz., eggs, larvae and pupae were studied at each of seven constant temperatures viz., 15 ± 1 , 18 ± 1 , 21 ± 1 , 24 ± 1 , 27 ± 1 , 30 ± 1 and $33 \pm 1^\circ\text{C}$ in BOD incubators. Oviposition periods of females were recorded in five replicates at each of the temperatures. Likewise, freshly laid eggs were incubated at different constant temperatures in five replicates and observed for hatching between 11.00 AM-12.30 PM daily. Similarly, 10 neonate larvae were reared at different temperatures and observed for moulting, mortality and pupation every day. Similarly, pupae were observed for adult emergence. Incubation period of eggs, and development periods of different larval instars and pupae were recorded at each of the temperatures.

Computation of thermal constants and developmental thresholds

According to rule of the constant sum of effective temperatures

$$(T-T_0) \times D = K \quad (\text{Davidson 1944, Wigglesworth 1972}).$$

Where T = temperature at which insect species is reared; T_0 = lower development threshold; D = duration of development, and K = thermal constant.

Development rates (r) of different developmental stages of *S. litura* were computed as the reciprocal of developmental periods in days (1/D) at each of the temperatures. The relationship between T and r was determined using linear regression where $r = a + bT$. The K was estimated as reciprocal of regression coefficient (b) between development rate and temperature.

$$K = 1/b \quad (\text{Kipyatkov and Lopatina 2010})$$

The T_0 was determined as ratio of regression intercept (a) and b.

$$T_0 = -a/b$$

Thermal constant and lower developmental thresholds were determined for eggs, larvae and pupae of *S. litura*.

RESULTS AND DISCUSSION

Oviposition by *S. litura* was observed only at 27°C and 33°C and at other constant temperatures either adults did not emerge or they died before oviposition. Neonate larvae completed their development from 1st instar to pupal stage at $18-33^\circ\text{C}$ (Table 1). However, larval development was not completed at 15°C as larvae died in 5th instar at 15°C . This might be due to the fact that at constant temperatures of 15°C , larvae required considerably more time to develop and favorable physiological conditions could not be maintained for such a long time. Developmental periods of different *S. litura* instars declined with an increase in temperature from 15 to 33°C . However, at 15°C larvae did not complete their life cycle. Development period of pupae also declined with temperature increase from 18°C to 33°C . Combined period for the larval and pupal development was 101.67 ± 0.67 days and 30 ± 1 days at 18 and 33°C , respectively (Table 3). The larval and pupal development periods of *S. litura* were thus influenced by temperature, which has also been observed earlier (Prasad 2004; Bharathi *et al.*, 2007; Patil *et al.*, 2014). Each species has an ecological maxima and minima within

Table 1: Developmental period in days (mean±SE) of *S. litura* at different constant temperatures

Temp.	n	I instar	II instar	III instar	IV instar	V instar	VI instar	First instar to final instar	Pupal	Total period
15°C	10	22.50±0.65	21.14±1.70	18.33±3.15	19.33±3.38	-	-	-	-	-
18°C	10	20.33±1.20	19.56±0.75	11.38±1.03	10.83±0.98	6.00±0.12	8.67±0.33	69.67±1.20	32.00±1.53	101.67±0.67
21°C	10	3.22±0.15	7.00±0.50	10.40±0.93	6.60±0.93	6.40±0.60	8.25±1.03	41.25±1.18	20.25±0.95	61.50±0.65
24°C	10	4.89±0.31	6.89±0.45	8.50±0.76	4.50±0.71	4.50±0.19	4.88±0.35	33.67±0.92	12.14±0.51	45.83±0.79
27°C	10	4.63±0.32	5.25±0.52	6.80±1.36	4.60±0.81	3.00±0.55	5.75±0.75	30.00±2.52	13.67±0.33	43.67±2.85
30°C	10	5.00±0.31	5.43±0.65	7.71±0.29	5.83±0.31	3.83±0.40	5.17±0.31	33.17±0.79	11.00±1.15	45.33±2.03
33°C	10	3.30±0.15	4.20±0.97	4.00±0.58	3.00±0.00	2.33±0.33	4.33±0.67	22.00±1.00	8.00±1.00	30.00±1.00
CD (0.05)		1.512	3.344	3.251	2.139	0.931	1.088	2.288	1.623	2.438

Table 2: Larval and pupal survival (%) in *S. litura* at different constant temperatures

Temp.	n	I instar	II instar	III instar	IV instar	V instar	VI instar	Total larval survivorship	Pupal survivorship	Total survivorship
15°C	10	80.00	87.50	85.72	50.00	0.0	0.0	0.0	0.0	0.0
18°C	10	90.00	100.00	88.89	75.00	83.33	60.00	30.00	30.00	30.00
21°C	10	90.00	100.00	55.56	100.00	100.00	80.00	40.00	40.00	40.00
24°C	10	90.00	100.00	88.89	100.00	100.00	100.00	80.00	70.00	60.00
27°C	10	80.00	100.00	62.50	100.00	100.00	80.00	40.00	30.00	30.00
30°C	10	70.00	100.00	100.00	85.72	100.00	100.00	60.00	30.00	30.00
33°C	10	100.00	50.00	60.00	100.00	100.00	100.00	30.00	30.00	30.00

which its development rate increases with rise in temperature but decreases beyond it. Generally, lower temperatures result in a decrease in the rate of development and an increase in duration of a developmental stage (Manimanjari *et al.*, 2014). Present study revealed that *S. litura* could develop under constant temperatures ranging between 20°C and 30°C. We also found that the females of *S. litura* were able to lay eggs only at constant temperatures of 27°C and 33°C. Present results are in agreement with those reported by Rao *et al.* (1989), however, with a deviation that they reported egg laying at low temperature of 15°C, which we did not observe. Rao *et al.* (2014) reported decrease in development time of different stages with an increase in temperature from 20-35°C that was similar to the decrease in developmental time from 15-33°C in the present study too. Likewise, thermal requirement from first to sixth instar in present study occurred to be 527DD in comparison to 538.5 DD from egg to adult development in case of Rao *et al.* (2014).

Incubation, larval and pupal developmental periods of *S. litura* as recorded in present study were proximal to those reported earlier under similar temperature conditions (Fand *et al.*, 2015; Nandita *et al.*, 2017). In current study, *S. litura* completed its larval development in 6 instars that has also been reported earlier (Rao *et al.* 1989). The developmental durations of *S. litura* immature stages observed at different constant temperatures were largely in conformity with earlier reports (Rao *et al.*, 1989; Patil *et al.*, 2014). Deviations in the developmental periods in the present study vis-à-vis earlier work may be attributed to the different host plants used as food for rearing the insect species (Xue *et al.*, 2010).

In present study, the larval survival was observed to be maximum (80%) at 27°C and minimum (30%) at both 18 and 33 °C as indicated by pupal formation. Overall, survival from 1st instar to adult emergence ranged between a maximum

Table 3: Linear regression equation, lower developmental thresholds and thermal constants of larval and pupal developmental stage of *S. litura*.

Life stages	b	a	DD	To	R ²
I instar	0.013	-0.167	76.92	12.85	0.900
II instar	0.011	-0.129	90.90	11.73	0.958
III instar	0.008	-0.079	125	9.88	0.794
IV instar	0.012	-0.124	83.33	10.33	0.783
V instar	0.016	-0.160	62.5	10.00	0.785
VI instar	0.011	-0.120	90.91	10.91	0.869
I-VI instar	0.001	-0.013	1000	13.00	0.848
Pupa	0.005	-0.066	200	13.20	0.904

of 60% at 24°C and 7% at each of 18°C and 33°C (Table 2). The 24°C was thus observed to be the optimum temperature for the survival and development of *S. litura*. The survival rates of *S. litura* immature stages as well adult stage varied significantly at various constant temperatures. The constant temperatures below 20°C and above 35°C were highly unfavourable for survival of all the life stages, at which increased mortality rates were observed. Earlier, Fand *et al.* (2015) observed highest percentage of survival (65.0%) of *S. litura* larva when reared at constant temperature of 30°C in laboratory conditions. Manimanjari *et al.* (2014) reported the similar trend in survivorship of *S. litura* where in they found lowest and highest survival at 20°C and 30°C temperatures, respectively, when reared on sunflower (*Helianthus annuus* L.). Further, Xue *et al.* (2010) reported that survival rates of immature stages of *S. litura* are highly dependent on host plants used for larval feeding.

Based on regression of developmental rates on corresponding temperatures, thermal constant for 1st to 6th larval instars and pupal stage of *S. litura* were computed as 76.9, 90.9, 125, 83.3, 62.5, 90.9 and 200 degree days (DD), respectively, with corresponding development thresholds as 12.8, 11.7, 9.8, 10.3, 10, 10.9 and 13.2°C (Table 3; Fig 1). In present study, developmental threshold for different larval instars reared on artificial diet ranged from 9.8-13.2°C and single developmental threshold for larval stage was determined as 13°C. The lower developmental threshold temperatures for *S. litura* larval stages estimated in the present study were in closer agreement with the earlier reports (Rao, 1989). The slight deviations that exist between current thresholds and those reported earlier may be ascribed to different rearing conditions and host plants used as larval food. In present study, thermal constant for pupal stage was computed as 200

DD with developmental threshold as 13.2°C. Earlier Fand *et al.* (2015) reported lower developmental threshold for pupal stage of *S. litura* as 9.8°C.

It has also been opined earlier that variable values obtained for thermal constants in different environments probably are indicative of different ecotypes of the pest. Fecundity and adult longevity in *S. litura* are influenced by temperature, humidity, and larval and adult nutrition (Zhu *et al.*, 2000; Xue *et al.*, 2010). In present study, although larvae did not complete their development at 15 and 18°C, yet developmental thresholds for different development stages ranged from 9.8–13.2°C. This was possible due to extrapolation of developmental rates obtained for temperatures from 18-33°C.

In this study, thermal constants for different larval instars ranged from 62.5-90.9 DD, while their developmental thresholds ranged between 9.8-12.8°C. On the other hand, pupal development required more amount of degree days for its completion with higher developmental threshold. Pupa being a transformational stage in insect development thus required higher amount of heat energy and its development was also triggered at higher developmental threshold.

CONCLUSION

The growth and development of *S. litura* were significantly influenced by temperature. Results from this study showed that both low and high temperatures limited the survival and development of this insect pest and based on development, survival and fecundity, optimum temperature for the larval and pupal survival was observed to be around 24°C, though the development rates increased with temperature up to 30°C. On the other hand, optimum temperature ranges for oviposition seems to be 27-33°C. Data




Fig. 1: Linear relationship between the temperature and developmental rate of different developmental stages of *S.litura*.

on thermal constant and developmental thresholds of different developmental stages of *S. litura* can be utilized to formulate a population simulation model of the pest. Different abiotic and biotic mortality factors can also be incorporated in to the simulation model. Such a model will be useful to investigate the impact of global warming on biological processes that are responsible for changes in population levels of the pest. The model will thus prove more useful than empirical

insect-weather regression models that do not explain the mechanisms behind insect population change.

REFERENCES

- Bale, J.S., Masters, G.J., Hodkinson, I.D., Awmack, C., Bezemer, T.M., Brown, V.K., Butterfield, J., Buse, A., Coulson, J.C., Farrar, J. and Good, J.E.(2002). Herbivory in global climate change research: direct

- effects of rising temperature on insect herbivores. *Global Change Biol.*, 8: 1–16.
- Bharathi, J.L., Sriedhar, U., Kishore, B., Prasad, J.V. and Prasad Rao, J.A.V. (2007). Life table studies of *Spodoptera litura* on different types of tobacco. *Tobacco Res.*, 33: 36–42.
- CABI (2020). Invasive Species Compendium: Datasheets, maps, images, abstracts and full text on invasive species of the world. Available: <http://www.cabi.org/isc/datasheet/44520>. Accessed 11 June 2020.
- Duraimurugan, P. (2018). Effect of weather parameters on the seasonal dynamics of tobacco caterpillar, *Spodoptera litura* (Lepidoptera: Noctuidae) in castor in Telangana State. *J. Agrometeorol.*, 20(2): 139-143.
- EPPO (2020) .PQR database: European and Mediterranean Plant Protection Organization, Paris, France. Available: <http://www.eppo.int/DATABASES/pqr/pqr.htm>.
- IPCC. (2013). Climate Change 2013: The Physical Science Basis. In: Stocker TF, Qin D, Plattner GK, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM, editors. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge (UK) and New York (USA): Cambridge University Press. p. 1535.
- Jha, G. K., Singh, G., Vennila, S., Rao, M. S., Panwar, H. and Hegde, M. (2016). Predicting the severity of *Spodoptera litura* on groundnut in relation to climatic variability using ordinal logistic model. *J. Agrometeorol.*, 18(2): 282.
- Kroschel, J., Sporleder, M., Tonnang, H.E.Z., Juarez, H., Carhuapoma, P., Gonzales, J.C. and Simon, R. (2013). Predicting climate-change-caused changes in global temperature on potato tuber moth *Phthorimaea operculella* (Zeller) distribution and abundance using phenology modeling and GIS mapping. *Agric For Meteorol* 15: 228–241.
- Manimanjari, D., Rao, M.S., Swathi, P., Rao, C.A.R., Vanaja, M. and Maheswari, M. (2014). Temperature- and CO₂-Dependent Life Table Parameters of *Spodoptera litura* (Noctuidae: Lepidoptera) on Sunflower and Prediction of Pest Scenarios. *J Insect Sci.*, 14(1):297
- Nandita, T., Gotyal, B.S., Selvaraj, K. and Satpathy, S. (2017). Effect on biology of jute indigo caterpillar, *Spodoptera litura* (Fabricius) under five different constant temperatures. *J. Entomol. Zool. Studies* 5(4):102-106
- Patil, R.A., Mehta, D.M. and Jat, B.L. (2014). Studies on life fecundity tables of *Spodoptera litura* Fabricius on tobacco *Nicotiana tabacum* Linnaeus. *Entomol Ornithol Herpetol.*, 3: 1–5.
- Prasad, J.V. (2004). Comparative life table studies of *Spodoptera litura* Fabricius and *Spodoptera exigua* on FCV tobacco seedlings under southern light soil conditions. *Tobacco Res.*, 30: 42–47.
- Rao, C.V.R., Wightman, J.A. and Rao, D.V.R. (1989). Threshold temperatures and thermal requirements for the development of *Spodoptera litura* (Lepidoptera: Noctuidae). *Environ Entomol.*, 18: 548–551.
- Rao, M. S., Manimanjari, D., Rao, A. C. R., Swathi, P. and Maheswari, M. (2014). Effect of climate change on *Spodoptera litura* Fab. on peanut: a life table approach. *Crop Prot.*, 66: 98-106.
- Xue, M., Pang, Y.H., Wang, H.T, Li, Q.L. and Liu, T.X. (2010). Effects of four host plants on biology and food utilization of the cutworm, *Spodoptera litura*. *J Insect Sci.*, 10: 22.
- Zhu, S., Lu, Z., Chen, L., Yu, W. and Zhang S. (2000). Effect of temperature and food on *S. litura* population. *The J Appl Ecol.*, 11(1):111-114.